

Late Life Anxiety and Depression Symptoms, and Suicidal Behaviors in Racial/Ethnic Minority Older Adults in Community-Based Organizations and Community Clinics in the U.S.

Aida L. Jiménez¹, Mario Cruz-Gonzalez², Tiara Forsyth Calhoun², Lauren Cohen², and Margarita Alegría^{2, 3}

¹ Department of Psychology, University of Puerto Rico

² Disparities Research Unit, Department of Medicine, Massachusetts General Hospital, Boston, Massachusetts, United States

³ Departments of Medicine and Psychiatry, Harvard Medical School

Objective: Late life anxiety and depression represent a significant source of disability, with racial/ethnic minority older adults in the U.S. showing marked disparities in healthy aging. Community-based organizations (CBOs) and community clinics serve these populations for preventive care, yet few identify their mental health service needs. We examine the association between race/ethnicity and risk of mild-to-severe symptoms of anxiety and depression, and suicidal behaviors in minority older adults. **Method:** Data come from the multisite randomized controlled trial *Building Community Capacity for Disability Prevention for Minority Elders*, which screened 1,057 adults (45.5% Asian, 26.8% Latinx, 15.0% non-Latinx Black, 8.5% non-Latinx White, and 4.2% American Indian) aged 60+ years at CBOs and clinics in Massachusetts, New York, Florida, and Puerto Rico. Screened participants completed the Generalized Anxiety Disorder-7 (GAD-7) for anxiety symptoms, the Geriatric Depression Scale-15 (GDS-15) for depression symptoms, and the Paykel Suicide Risk Questionnaire for suicidal behaviors. **Results:** 28.1% of older adults reported mild-to-severe anxiety symptoms, 30.1% reported mild-to-severe depression symptoms, and 4.3% reported at least one suicidal behavior. Compared to non-Latinx Whites, Latinxs had higher odds of mild-to-severe anxiety and depression symptoms and one or more suicidal behaviors, and Asians had higher odds of mild-to-severe depression symptoms only. **Conclusions:** There is an urgent need to improve outreach for screening and preventive mental health care for minority older adults. Expanding outreach and community-based capacity to identify and treat minority older adults with mental health conditions represents an opportunity to prevent disability.

Public Significance Statement

Our results underscore the importance of outreaching minority older adults participating in community-based organizations (CBOs) and clinics for mental health screening given the high prevalence of anxiety and depressive symptoms, and suicidal behaviors. CBOs and clinics may offer opportunities for early identification of their mental health problems and provide mental health literacy and connection to secondary prevention programs to mitigate their higher risk of disability.

Keywords: racial/ethnic minority older adults, anxiety, depression, suicidal behaviors

This article was published Online First February 3, 2022.

Aida L. Jiménez  <https://orcid.org/0000-0002-1614-9390>

Mario Cruz-Gonzalez  <https://orcid.org/0000-0002-9888-3205>

Lauren Cohen  <https://orcid.org/0000-0002-5787-699X>

This study was funded by Grant 5R01AG046149-04 from the National Institute of Aging and National Institute of Mental Health (PI: Margarita Alegría). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The funders (NIA, NIMH) had no role in design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

The authors have no conflicts of interest to disclose.

Aida L. Jiménez played a lead role in conceptualization and writing of original draft and an equal role in investigation, methodology, supervision and writing of review and editing. Mario Cruz-Gonzalez played a lead role in

data curation, formal analysis, and software and an equal role in methodology, writing of original draft, and writing of review and editing. Tiara Forsyth Calhoun played a supporting role in writing of original draft and writing of review and editing. Lauren Cohen played a supporting role in writing of review and editing. Margarita Alegría played a lead role in conceptualization, funding acquisition, investigation, methodology, supervision, and visualization and an equal role in writing of original draft and writing of review and editing.

The authors would like to thank the participants, research staff, and partnering community organizations for their contributions to the study, and Sheri Lapatin Markle, Larimar Fuentes, and Isabel O'Malley for their feedback and revisions to the manuscript.

Correspondence concerning this article should be addressed to Aida L. Jiménez, Department of Psychology, University of Puerto Rico, Río Piedras Campus, P.O. Box 23345, San Juan, PR 00931-3345, United States. Email: Aida.jimenez@upr.edu

The U.S. Census Bureau projects that by 2060 nearly one in four people in the U.S. will be ages 65+ (Medina et al., 2020). Many U.S. older adults have low income or belong to diverse racial/ethnic groups (23%), and experience higher and earlier morbidity and mortality than non-Latinx Whites (Davison et al., 2019; Gerst et al., 2010). These changing demographics will result in a disproportionate rise in age-related health conditions associated with disability, such as elevated anxiety and depression symptoms, leading to substantial morbidity (Jones, 2012; Morin et al., 2020; Yang & Levkoff, 2005).

About 12.6% of U.S. community-dwelling adults ages 65+ suffer from anxiety disorders (Patel et al., 2019), and 13.7% of adults ages 55+ meet criteria for major depressive disorder (Laborde-Lahoz et al., 2015). Although prevalence of these disorders declines with age (Byers et al., 2010; Hybels & Blazer, 2003), late life mental health symptoms remain common, particularly in women. Despite effective diagnostic tools and therapies for late life anxiety and depression, these disorders remain undiagnosed and untreated (Grenier et al., 2019; Knight, 2019) which is associated with increased morbidity and mortality, decreased quality of life, increased cost of care, and higher risk of suicide (Cullen, 2019; Feng et al., 2013; Ho et al., 2014). Older adults are also at higher risk of suicide than middle aged adults due to physical and cognitive decline and cumulative stressors (Almeida et al., 2012; Bennardi et al., 2019; Chang et al., 2017; Fässberg et al., 2016; Lutz & Fiske, 2018). Late life suicidal behaviors are often confounded with other health conditions, psychological distress, or dementia, making it difficult to detect (Hegeman et al., 2012; Van Orden & Conwell, 2016).

Late life anxiety and depression can be undetected and untreated due to a misperception that mental health decline is a normal part of aging (Balsamo et al., 2018; Kiosses & Marino, 2019; Knight, 2019;). In addition, racial/ethnic minority older adults encounter increased stigma around mental health-care (Jimenez et al., 2013) and are exposed to cumulative stressors associated with higher risk of poor mental health (Casado & Leung, 2002; Kang et al., 2013; Sadavoy et al., 2004). Evidence suggests that minority older adults may be at higher risk of poor mental health because of cumulative stressors, such as economic inequity, differences in cultural values, immigration status, and acculturation (Aranda et al., 2001), language barriers (Chen et al., 2009; Derr, 2016; Kim et al., 2015; Mui & Kang, 2006), and lack of access to healthcare (Aranda et al., 2001).

Social Determinants of Health Framework

According to the World Health Organization (WHO, 2014), mental health is shaped by social, cultural, and economic factors, and physical environments. Social determinants of health (SDH) such as income, education, employment, gender, age, environment, cultural and social values have been associated with mental disorders (Lund et al., 2018; Solar & Irwin, 2010; WHO, 2021). We draw from the SDH framework to understand factors associated with anxiety, depression, and suicidal behaviors in minority older adults.

Economic Inequity

Prior work has shown that poverty and financial strain, as SDH, may contribute to increased risk of depression in racial/ethnic minority older adults compared to non-Latinx Whites (Angel

et al., 2003; García et al., 2019; Kahn & Fazio, 2005). Economic difficulties can expose these minorities to additional stressors (e.g., lack of food, medicine, and utilities), further increasing their vulnerability to depressive symptoms (Chiriboga et al., 2002; Krause, 1987; Krause et al., 1991). Older Latinx are more likely to live below the federal poverty level (Administration for Community Living, 2018) and to experience financial stress associated with greater depressive symptoms (Chiriboga et al., 2002). Low socioeconomic status among African Americans is also associated with increased risk of depressive symptoms (Bailey et al., 2019; Holden et al., 2013).

Differences in Cultural Values

Culture, as a broad construct including shared attitudes, beliefs, and norms, might relate to how diverse groups define what is “normal” and shape how they explain, experience, and respond to mental distress, including help-seeking behaviors (Torres et al., 2020). Similarly, “ethnicity,” defined as common ancestry and shared cultural traditions specific to social context and heritage (Cokley & Moore, 2007), can influence interpretation of mental health symptoms.

Immigration Status and Acculturation

Immigration status and low levels of acculturation are also associated with increased risk of depression in the U.S. (Aranda et al., 2001). Low acculturation can be a stressor among Latinx immigrants (Caplan, 2007), with longer duration in the U.S. correlating with worse mental health (Perreira et al., 2015). Immigrant older adults often face a lifetime of adjustment, which can contribute to poor mental health (Lum & Vanderkaa, 2010). Gerst et al. (2010) found that older Mexican American immigrants were at higher risk for depressive symptoms compared to U.S.-born Mexican Americans, a finding that has been replicated in Asian older adults (Mui & Kang, 2006). Low levels of acculturation might also lead to family conflict due to dissonance among family members of different generations (Lau et al., 2005). Older Latinx may also expect their children to care for them and follow their cultural values of “familismo” and respect (Jimenez et al., 2019). Limited English proficiency and language preference, often used as proxies of acculturation, are also linked with less access to medical care and poor health status and mental health among ethnic minorities, including Latinx, Asians, and African Americans (DuBard & Gizlice, 2008; Kim et al., 2011; Sentell et al., 2007; Wisnivesky et al., 2012).

Access to Healthcare

Mental health treatment for underserved racial/ethnic minority older adults is mostly centered in primary care (Jimenez et al., 2012; Snowden & Pingitore, 2002). Yet, primary care providers (physicians, nurse practitioners, and physician assistants) and behavioral health care providers (psychiatrists, psychologists, and social workers) often do not detect or treat mental disorders among ethnic minorities (Jones et al., 2018; Tai-Seale et al., 2007; Ward et al., 2016) given their lack of training in cultural differences and unawareness of how anxiety and depression symptoms might manifest (Bailey et al., 2019).

Current Gap in Knowledge

Research done in primary care elucidates intervention trends and targets for anxiety, depression, and suicidal behaviors among older adults (Kroenke et al., 2007; Luoma et al., 2002; Park & Unützer, 2011). However, many racial/ethnic minority older adults may not seek services due to lack of trust and prior negative experiences, as well as linguistic barriers (Balakrishnan & Jordan, 2019; Glover et al., 2017; Hansen et al., 2016; Musa et al., 2009). Community-based organizations (CBOs), where these adults congregate, offer a unique opportunity to identify those with unmet need and serve them. CBOs are important health system stakeholders providing services to the most marginalized members of their community (Blas et al., 2008; Chillag et al., 2002) encompassing a range of organizations, including councils on aging, senior and day centers, faith-based organizations, and community health centers (Ramanadhan et al., 2012). Most CBOs offer services to maintain patients' health and well-being, such as educational programs, legal assistance, nutritional support, transportation, housing, and financial services (Sieglar et al., 2015; Turner et al., 2019; Wu et al., 2019). In contrast, community clinics provide affordable medical care. CBOs and clinics can aid in promoting mental health literacy, social inclusion, mental health awareness, and reducing stigma around receipt of mental health services (Evans-Lacko et al., 2014; Jorm, 2012; Turner et al., 2019; Wu et al., 2019). However, little is known about the prevalence and correlates of mental health symptoms among racial/ethnic minority older adults who receive services through CBOs and clinics.

The Present Study

Our study addresses the current gap in knowledge regarding rates of clinically significant anxiety and depression symptoms and suicidal behaviors in racial/ethnic minority older adults receiving services in CBOs and clinics across four U.S. states (Massachusetts, New York, Florida, and Puerto Rico). Since poor mental health for racial/ethnic minority older adults is associated with barriers including economic inequity (Bailey et al., 2019; Chiriboga et al., 2002; Holden et al., 2013; Krause, 1987; Krause et al., 1991), immigrant status (Aranda et al., 2001), and English language proficiency (Kim et al., 2011), we sought to examine racial/ethnic differences in mental health outcomes in a community-based sample of low-income mostly immigrant minority older adults. This contextual approach provides an opportunity to disentangle overlapping contributing factors to poor mental health in these minorities and highlights the importance of mental health screening within the CBOs and clinics where they seek care. Specifically, we (a) describe demographic and mental health outcomes of minority older adults screened as part of a larger randomized controlled trial; (b) explore the distribution of these outcomes by place of recruitment (CBOs vs. clinics); (c) evaluate the association between race/ethnicity and mild-to-severe anxiety and depression symptoms and any suicidal behavior; and (d) examine the association between preferred language and mental health outcomes. We hypothesized that racial/ethnic minority older adults would have increased odds of mild-to-severe anxiety and depression symptoms and any suicidal behavior compared to non-Latinx Whites (Alegría, Frontera, et al., 2019). Secondary hypotheses were that we would observe higher prevalence of depression among older adults recruited at clinics compared to CBOs (Ell, 2006); that adults ages 75+ across racial/ethnic groups

would report better mental health outcomes than adults ages 60-74 (Hybels & Blazer, 2003); and that non-English speakers would be at increased risk of poor mental health than English-speakers (Kim et al., 2019).

Method

Sampling and Procedure

We used data from the multisite randomized controlled trial *Building Community Capacity for Disability Prevention for Minority Elders* (Alegría, Frontera, et al., 2019), which combined mental health and physical disability prevention interventions in nine CBOs and two community-based clinics in four U.S. states: Massachusetts (2 clinics and 5 CBOs), New York (2 CBOs), Florida (1 CBO), and Puerto Rico (1 CBO). Data were collected between May 2015 and March 2019, with 1,057 participants screened for eligibility. Screening was administered only if participants were 60+ years and fluent in English, Spanish, Mandarin, or Cantonese. Research assistants (RAs) approached potential participants at the collaborating CBOs and clinics to describe the screening process or conducted outreach events at participating sites. If the participant showed interest, the RA arranged an in-person meeting for informed consent using the University of California, San Diego (UCSD) Brief Assessment of Capacity to Consent (Jeste et al., 2007). Participants who passed capacity to consent were screened for eligibility through an in-person interview in their preferred language.

In the present study, we used data from the 1,057 participants screened for eligibility, from whom 307 were eligible and agreed to participate in the trial (Alegría, Frontera, et al., 2019). These 1,057 participants were compensated with a \$10 gift card after completing the screening. Screening data included demographic information and an assessment of mental and physical health symptoms described below. The protocol was approved by the Institutional Review Boards of Massachusetts General Hospital/Mass General Brigham (protocol number 2015P001505) and New York University (protocol 114-01903), with ceded reviews for CBOs and clinics conducting human subject's research.

Outcome Measures

Mild-to-Severe Anxiety Symptoms

Anxiety symptoms were assessed via the Generalized Anxiety Disorder-7 (GAD-7), a 7-item self-report screening of symptom severity of generalized anxiety disorder (GAD; Spitzer et al., 2006). Respondents are asked how often, in the last 2 weeks, they have been bothered by each of the core GAD symptoms (0 = *not at all* and 3 = *nearly every day*). The scale ranges from 0 to 21, with higher scores reflecting worse anxiety symptoms. Using a threshold score of 10, the GAD-7 has a sensitivity of 89% and a specificity of 82% for GAD. Scores of five and above indicate *mild-to-severe anxiety symptoms* (Spitzer et al., 2006); thus, our outcome variable was an indicator equal to one if the GAD-7 ≥ 5 and equal to zero otherwise. Internal consistency (Cronbach's α) was .89 in the overall sample, and .86, .86, .92, and .93 in English, Spanish, Mandarin, and Cantonese, respectively.

Mild-to-Severe Depression Symptoms

Depression symptoms were assessed using the Geriatric Depression Scale-15 (GDS-15), a 15-item self-report screening of clinical depression among older adults (Sheikh & Yesavage, 1986). Participants are asked *yes* or *no* questions on how they felt the previous week. The scale ranges from 0 to 15, with higher scores reflecting more depression symptoms. A score of five and above is indicative of *mild-to-severe depression symptoms* (Sheikh & Yesavage, 1986); thus, our outcome variable was an indicator equal to one if the GDS-15 ≥ 5 and equal to zero otherwise ($\alpha = .84$ in the overall sample, and .79, .84, .86, and .87 in English, Spanish, Mandarin, and Cantonese, respectively).

Suicidal Behaviors

Suicidal behaviors were assessed using the Paykel Suicide Risk Questionnaire, a 5-item self-report screening (*yes/no* questions) for assessing passive and active suicidal ideation and behaviors within the past month (Paykel et al., 1974). Our outcome variable was an indicator equal to one if the participant endorsed any of the Paykel items (*yes* to either question) and equal to zero otherwise. We refer to this indicator as *any suicidal behavior*.

Independent Variables

Race/Ethnicity

Race and ethnicity were assessed using self-reported information. Participants were first classified as Latinx if they self-identified as of Spanish, Hispanic, or Afro-Caribbean descent ($N = 283$). The remaining participants were classified as Asian if they self-identified as Asian/Pacific Islander ($N = 481$); as American Indian ($N = 44$); as non-Latinx Black if they self-identified as Black/African/African American ($N = 159$); and as non-Latinx White if they self-identified as White but not as of Latinx descent ($N = 90$).

Interview Language

Preferred interview language was used as a proxy for acculturation, and was assessed through the question *Would you be comfortable participating in a program in any of the following languages? If so, which one?* Participants could select more than one language. Response options included *English, Spanish, Mandarin, Cantonese, and none of these*. These language options were provided, because participants were eligible for the trial only if they spoke at least one of them. When participants selected more than one language ($N = 119$), they indicated the one that they preferred to be interviewed.

Previous literature has documented measurement invariance by race/ethnicity and language for the GAD-7, which has been found to have the same factor structure and equivalent response patterns among Latinx Americans with English or Spanish language preferences (Mills et al., 2014). No studies assessing the measurement properties of the GDS-15 or the Paykel Suicide Risk Questionnaire were identified (Cassidy et al., 2018). We conducted our analyses with the potential limitation that observed outcome differences might reflect differences in interpretation for some of our outcome measures.

Additional Covariates

Known correlates of our three outcomes were included as covariates (Aranda et al., 2001; Byers et al., 2010; Chiriboga et al., 2002; Hong et al., 2014; Richardson et al., 2011).

Socio-Demographics

We included age (60–64, 65–74, and 75+ years), gender (male, female), and education (less than high school, high school diploma, or more).

Physical Health

Physical health was measured using the Short Physical Performance Battery (SPPB), an objective tool from the National Institute of Aging to evaluate lower extremity functioning in older persons (Guralnik et al., 1994). It includes three tests of functional relevance: standing balance, 4-m gait speed, and five-repetition sit-to-stand motion. Each test is scored on a 5-point scale from 0 (unable to perform/low performance) to 4 (high performance) and summed for a score range between 0 and 12, with higher scores representing better physical functioning. SPPB scores were used as a continuous measure.

Site of Recruitment

A binary variable equal to one if participants were recruited in CBOs and equal to zero if they were recruited in clinics was also included, because place of recruitment could be associated with the outcomes.

Data Analysis

Descriptive Statistics

We described the outcome measures and covariates in the total sample and by race/ethnicity, comparing non-Latinx Whites and racial/ethnic minority groups (Table 1). Statistical comparisons used cluster-robust standard errors to account for nonindependence of observations within our 11 sites (CBOs + clinics). Thus, the joint test of equality across racial/ethnic groups had an $F(h, G - 1)$ distribution, where h is the number of (linearly independent) restrictions, and G is the number of clusters (CBOs + clinics). We also compared the distribution of the outcomes by site of recruitment (clinics vs. CBOs; Table 2). Statistical comparisons between clinics and CBOs also used cluster-robust standard errors, so the joint test of equality across clinics and CBOs had the same $F(h, G - 1)$ distribution.

Association Between Race/Ethnicity and Outcome Measures

We used logistic regression models to estimate the association between race/ethnicity and *mild-to-severe anxiety symptoms* (GAD-7 ≥ 5), *mild-to-severe depression symptoms* (GDS-15 ≥ 5), and *any suicidal behavior* (endorsing any Paykel item), controlling for the covariates of gender, education, age, place of recruitment, and physical health (Table 3). To account for nonindependence among participants from the same CBOs or

Table 1
Patient Descriptive Statistics in the Overall Sample and by Race/Ethnicity

Variable	Total screened (<i>N</i> = 1,057) <i>N</i> (%)	Non-Latino White (<i>N</i> = 90) <i>N</i> (%)	Non-Latino Black (<i>N</i> = 159) <i>N</i> (%)	American Indian (<i>N</i> = 44) <i>N</i> (%)	Asian (<i>N</i> = 481) <i>N</i> (%)	Total screened (<i>N</i> = 1,057) <i>N</i> (%)	<i>F</i> (<i>h</i> , <i>G</i> - 1); ^a <i>p</i>
Outcomes							
Anxiety symptoms ^b							
None or minimal	760 (71.9%)	61 (67.8%)	111 (69.8%)	29 (65.9%)	391 (81.3%)	168 (59.4%)	<i>F</i> (4, 10) = 4.06; <i>p</i> = .03
Mild to severe	297 (28.1%)	29 (32.2%)	48 (30.2%)	15 (34.1%)	90 (18.7%)	115 (40.6%)	
Depression symptoms ^c							
None or minimal	671 (69.9%)	64 (77.1%)	105 (76.1%)	32 (76.2%)	293 (68.3%)	177 (66.0%)	<i>F</i> (4, 10) = 2.81; <i>p</i> = .08
Mild to severe	289 (30.1%)	19 (22.9%)	33 (23.9%)	10 (23.8%)	136 (31.7%)	91 (34.0%)	
Any suicidal behavior ^d							
No	1,012 (95.7%)	88 (97.8%)	153 (96.2%)	43 (97.7%)	465 (96.7%)	263 (92.9%)	<i>F</i> (4, 10) = 21.41; <i>p</i> < .01
Yes	45 (4.3%)	2 (2.2%)	6 (3.8%)	1 (2.3%)	16 (3.3%)	20 (7.1%)	
Demographics							
Gender							
Male	257 (24.3%)	14 (15.6%)	39 (24.5%)	16 (36.4%)	131 (27.2%)	57 (20.1%)	<i>F</i> (4, 10) = 5.31; <i>p</i> = .01
Female	800 (75.7%)	76 (84.4%)	120 (75.5%)	28 (63.6%)	350 (72.8%)	226 (79.9%)	
Education							
Less than high school	440 (41.7%)	14 (15.6%)	36 (22.8%)	23 (52.3%)	249 (51.8%)	118 (41.8%)	<i>F</i> (4, 10) = 12.72; <i>p</i> < .01
High school or more	615 (58.3%)	76 (84.4%)	122 (77.2%)	21 (47.7%)	232 (48.2%)	164 (58.2%)	
Interview language							
English	240 (22.7%)	84 (93.3%)	126 (79.2%)	10 (22.7%)	10 (2.1%)	10 (3.5%)	<i>F</i> (7, 10) = 1902.44; <i>p</i> < .01
Spanish	347 (32.8%)	5 (5.6%)	33 (20.8%)	34 (77.3%)	2 (4%)	273 (96.5%)	
Mandarin	199 (18.8%)	1 (1.1%)	0 (0%)	0 (0%)	198 (41.2%)	0 (0%)	
Cantonese	271 (25.6%)	0 (0%)	0 (0%)	0 (0%)	271 (56.3%)	0 (0%)	
Age							
60-64	60 (5.7%)	5 (5.6%)	5 (3.1%)	4 (9.1%)	10 (2.1%)	36 (12.7%)	<i>F</i> (8, 10) = 11.91; <i>p</i> < .01
65-74	418 (39.5%)	40 (44.4%)	73 (45.9%)	20 (45.5%)	157 (32.6%)	128 (45.2%)	
75+	579 (54.8%)	45 (50.0%)	81 (50.9%)	20 (45.5%)	314 (65.3%)	119 (42.0%)	
Place of recruitment							
Clinic	391 (37.0%)	75 (83.3%)	37 (23.3%)	18 (40.9%)	159 (33.1%)	102 (36.0%)	<i>F</i> (4, 10) = 8.79; <i>p</i> < .01
CBO	666 (63.0%)	15 (16.7%)	122 (76.7%)	26 (59.1%)	322 (66.9%)	181 (64.0%)	
Geographic location							
Massachusetts	749 (70.9%)	85 (94.4%)	60 (37.7%)	26 (59.1%)	374 (77.8%)	204 (72.1%)	<i>F</i> (8, 10) = 4.64; <i>p</i> = .01
New York	198 (18.7%)	2 (2.2%)	83 (52.2%)	5 (11.4%)	107 (22.2%)	1 (4%)	
Florida	11 (1.0%)	2 (2.2%)	0 (0%)	0 (0%)	0 (0%)	9 (3.2%)	
Puerto Rico	99 (9.4%)	1 (1.1%)	16 (10.1%)	13 (29.5%)	0 (0%)	69 (24.4%)	
Physical health (SPPB)							
<i>M</i> (<i>SD</i>)	7.88 (2.70)	7.15 (2.45)	7.17 (2.56)	7.27 (2.64)	8.65 (2.63)	7.32 (2.63)	<i>F</i> (4, 10) = 20.03; <i>p</i> < .01

Notes. CBO = community-based organization; SPPB = short physical performance battery.

^a Statistical comparison between non-Latino Whites and racial/ethnic minority groups used cluster-robust standard errors to account for the potential nonindependence of observations within our 11 CBOs and clinics. With cluster-robust standard errors, the joint test of equality across racial/ethnic groups has an *F*(*h*, *G*-1) distribution, where *h* is the number of (linearly independent) restrictions, and *G* is the number of clusters (11 CBOs and clinics). ^b Anxiety symptoms were assessed using the Generalized Anxiety Disorder-7 (GAD-7); scores of five and above are indicative of mild-to-severe anxiety symptoms (Spitzer et al., 2006). ^c Depressive symptoms were assessed using the Geriatric Depression Scale-15 (GDS-15); a score of five and above is indicative of mild-to-severe depressive symptoms (Sheikh & Yesavage, 1986). Ninety-seven patients were not administered the GDS-15. ^d Any suicidal behavior was assessed using the Paykel Suicide Risk Questionnaire.

Table 2*Distribution of Outcome Variables by Place of Recruitment*

Outcome	Total screened (<i>N</i> = 1,057) <i>N</i> (%)	Recruited in clinics (<i>N</i> = 391) <i>N</i> (%)	Recruited in CBOs (<i>N</i> = 666) <i>N</i> (%)	<i>F</i> (<i>h</i> , <i>G</i> - 1) ^a <i>p</i>
Anxiety symptoms ^b				
None or minimal	760 (71.9%)	266 (68.0%)	494 (74.2%)	<i>F</i> (1, 10) = 2.50; <i>p</i> = .14
Mild to severe	297 (28.1%)	125 (32.0%)	172 (25.8%)	
Depression symptoms ^c				
None or minimal	671 (69.9%)	263 (67.3%)	408 (71.7%)	<i>F</i> (1, 10) = 1.98; <i>p</i> = .19
Mild to severe	289 (30.1%)	128 (32.7%)	161 (28.3%)	
Any suicidal behavior ^d				
No	1,012 (95.7%)	375 (95.9%)	637 (95.6%)	<i>F</i> (1, 10) = .03; <i>p</i> = .86
Yes	45 (4.3%)	16 (4.1%)	29 (4.4%)	

Notes. CBO = community-based organization.

^aStatistical comparison between participants recruited in clinics and participants recruited in CBOs used cluster-robust standard errors to account for the potential nonindependence of observations within our 11 CBOs and clinics. With cluster-robust standard errors, the joint test of equality across clinics and CBOs has an *F*(*h*, *G*-1) distribution, where *h* is the number of (linearly independent) restrictions, and *G* is the number of clusters (11 CBOs and clinics). ^bAnxiety symptoms were assessed using the Generalized Anxiety Disorder-7 (GAD-7); scores of five and above are indicative of mild-to-severe anxiety symptoms (Spitzer et al., 2006). ^cDepressive symptoms were assessed using the Geriatric Depression Scale-15 (GDS-15); a score of five and above is indicative of mild-to-severe depressive symptoms (Sheikh & Yesavage, 1986). Ninety-seven patients were not administered the GDS-15. ^dAny suicidal behavior was assessed using the Paykel Suicide Risk Questionnaire.

clinics, our models used cluster-robust standard errors. Although we could have modeled nonindependence using multilevel models with a site-level random intercept, we did not, because the intraclass correlation coefficients were very low in our sample (between .006 and .055), and a likelihood ratio test versus a (single-level) logistic model could not reject the null hypothesis that the random intercepts were zero. Individual coefficients for categorical variables (i.e., race/ethnicity and age) were interpreted if the omnibus tests were significant at the .05

level. Odds ratios (*OR*) and their 95% Confidence Intervals (95% *CI*) were estimated using Stata 15 (Stata Corp, 2017).

Association Between Interview Language and Outcome Measures

Since interview language was highly correlated with race/ethnicity, our logistic models examining the association between race/

Table 3*Logistic Regression Model for the Association Between Race/Ethnicity and Mental Health Outcomes*

Independent variables	Dependent variable					
	Mild-to-severe anxiety symptoms		Mild-to-severe depression symptoms		Any suicidal behavior	
	(GAD-7 ≥ 5)		(GDS ≥ 5)		(Any Paykel item)	
	<i>OR</i> [95% <i>CI</i>] ^a	<i>P</i> > <i>z</i>	<i>OR</i> [95% <i>CI</i>] ^a	<i>P</i> > <i>z</i>	<i>OR</i> [95% <i>CI</i>] ^a	<i>P</i> > <i>z</i>
Gender (male reference)						
Female	1.41 [.94, 2.12]	.101	1.36 [.77, 2.39]	.284	1.40 [.79, 2.49]	.253
Race/Ethnicity (non-Latinx White reference)						
Non-Latinx Black	1.07 [.76, 1.52]	.693	1.35 [.84, 2.18]	.216	1.82 [.50, 6.67]	.363
American Indian	1.26 [.64, 2.45]	.504	1.17 [.67, 2.07]	.576	1.24 [.33, 4.66]	.755
Asian	.73 [.43, 1.23]	.238	2.62 [1.90, 3.61]	<.001	2.10 [.94, 4.68]	.069
Latinx	1.57 [1.13, 2.18]	.007	1.85 [1.39, 2.46]	<.001	3.59 [1.43, 8.99]	.006
Omnibus test	$\chi^2(4) = 13.43$	.009	$\chi^2(4) = 45.78$	<.001	$\chi^2(4) = 31.53$	<.001
Education (less than high school reference)						
High school or more	1.03 [.82, 1.29]	.822	.96 [.79, 1.18]	.715	1.60 [.90, 2.84]	.106
Age (60-64 reference)						
65-74	.69 [.41, 1.16]	.160	.56 [.35, .91]	.020	.59 [.37, .93]	.024
75+	.44 [.27, .74]	.002	.35 [.24, .51]	<.001	.52 [.31, .86]	.010
Omnibus test	$\chi^2(2) = 16.94$	<.001	$\chi^2(2) = 84.51$	<.001	$\chi^2(2) = 8.09$	.017
Place of recruitment (clinic reference)						
CBO	.78 [.67, .91]	.002	.78 [.63, .97]	.025	1.03 [.58, 1.82]	.928
Physical health: SPPB score	.89 [.82, .96]	.002	.86 [.78, .96]	.005	.93 [.89, .98]	.004
Observations	—	1,055	—	959	—	1,055

Notes. GAD-7 = Generalized Anxiety Disorder-7; GDS = Geriatric Depression Scale; *OR* = odds ratio; *CI* = confidence interval; CBO = community-based organization; SPPB = short physical performance battery.

^a95% confidence intervals were constructed using cluster-robust standard errors to account for the potential nonindependence of observations within our 11 CBOs and clinics.

ethnicity and the outcomes did not control for interview language. We applied this exclusion, because the effect of race/ethnicity could not be estimated independently from interview language (e.g., 99.5% and 100% of participants interviewed in Mandarin and Cantonese, respectively, self-identified as Asian; see Table 1). As a sensitivity analysis, we computed the same logistic models, but replaced race/ethnicity with interview language; that is, we estimated the association between interview language and the outcomes while controlling for gender, education, age, place of recruitment, and physical health (Table 4). These models also used cluster-robust standard errors, and we interpreted the significance of individual coefficients for interview language if the omnibus test was significant at the .05 level.

Results

Table 1 presents descriptive statistics for the total sample ($N = 1,057$). *Mild-to-severe anxiety* (GAD-7 ≥ 5) was reported by 28.1% of older adults, *mild-to-severe depression* (GDS-15 ≥ 5) by 30.1%, and *any suicidal behavior* per any Paykel item by 4.3%. Most respondents were female, had a high school diploma or higher degree, and ages 75+. Most participants were Asian, followed by Latinx, non-Latinx Blacks, non-Latinx Whites, and American Indians. Most respondents (63.0%) were recruited at CBOs, with the remainder recruited at clinics. Since the two clinics and five of the nine CBOs were in Massachusetts, most respondents were recruited in this state, followed by New York, Puerto Rico, and Florida.

Table 1 also presents descriptive statistics by race/ethnicity. More Latinx met criteria for *mild-to-severe anxiety* compared to non-Latinx Whites, while fewer Asians met these criteria. *Mild-to-severe*

depression was more prevalent for both Latinx and Asians compared to non-Latinx Whites, although differences across racial/ethnic groups were not significant ($p = .08$). More Latinx endorsed *any suicidal behavior* than non-Latinx Whites. Ninety-three percent of non-Latinx White and 79.2% of non-Latinx Black respondents were interviewed in English, while 97.5% of Asians responded either in Mandarin (41.2%) or in Cantonese (56.3%), and 96.5% of Latinx responded in Spanish. Most non-Latinx Blacks (76.7%), Asians (59.1%), and Latinx (66.9%) were recruited at CBOs, while more than 83% of non-Latinx Whites were recruited at clinics. Across racial/ethnic groups, most participants were recruited in Massachusetts except for non-Latinx Blacks, most of whom were recruited in New York. Table 1 also indicates that Asians had better physical health per the SPPB than non-Latinx Whites.

Table 2 shows the distribution of the outcome measures by place of recruitment. No significant differences were observed for either outcome.

Table 3 presents the results of logistic regression models for the association between race/ethnicity and our three outcomes. After controlling for covariates, significant differences across racial/ethnic groups were observed for all outcomes ($p < .05$ for all omnibus tests). Compared to non-Latinx Whites, Latinx had almost 1.6 higher odds of reporting *mild-to-severe anxiety symptoms* (OR [95% CI] = 1.57 [1.13, 2.18]), almost twice the odds of reporting *mild-to-severe depression symptoms* (OR [95% CI] = 1.85 [1.39, 2.46]), and more than three times the odds of reporting *any suicidal behavior* (OR [95% CI] = 3.59 [1.43, 8.99]). Asians had higher odds of *mild-to-severe depression symptoms* than non-Latinx Whites, but no differences were observed for *mild-to-severe anxiety symptoms* or *any suicidal behavior*. There were also no differences in the odds

Table 4

Logistic Regression Model for the Association Between Interview Language and Mental Health Outcomes

Independent variables	Dependent variable					
	Mild-to-severe anxiety symptoms		Mild-to-severe depression symptoms		Any suicidal behavior	
	(GAD-7 ≥ 5)		(GDS ≥ 5)		(Any Paykel item)	
	OR [95% CI] ^a	$P > z $	OR [95% CI] ^a	$P > z $	OR [95% CI] ^a	$P > z $
Gender (male reference)						
Female	1.43 [.95, 2.15]	.084	1.38 [.78, 2.45]	.275	1.46 [.85, 2.51]	.172
Interview language (English reference)						
Spanish	1.54 [1.02, 2.34]	.040	1.86 [1.20, 2.88]	.006	1.90 [.86, 4.18]	.111
Mandarin	.58 [.42, .80]	< .001	2.66 [1.79, 3.96]	< .001	1.74 [.50, 6.10]	.388
Cantonese	.79 [.54, 1.16]	.237	2.46 [1.71, 3.54]	< .001	1.05 [.32, 3.42]	.940
Omnibus test	$\chi^2(3) = 16.01$	.001	$\chi^2(3) = 39.63$	< .001	$\chi^2(3) = 8.27$	.041
Education (less than high school reference)						
High school or more	1.10 [.84, 1.43]	.492	.99 [.77, 1.27]	.934	1.45 [.83, 2.52]	.191
Age (60-64 reference)						
65-74	.67 [.40, 1.14]	.140	.59 [.35, .97]	.038	.56 [.31, 1.04]	.065
75+	.44 [.27, .74]	.002	.36 [.25, .53]	< .001	.46 [.23, .91]	.026
Omnibus test	$\chi^2(2) = 16.28$	< .001	$\chi^2(2) = 66.75$	< .001	$\chi^2(2) = 5.33$	.070
Place of Recruitment (clinic reference)						
CBO	.76 [.67, .86]	< .001	.79 [.61, 1.02]	.069	1.17 [.67, 2.06]	.578
Physical health: SPPB Score	.89 [.83, .96]	.003	.86 [.78, .96]	.007	.93 [.88, .98]	.005
Observations	—	1,055	—	959	—	1,055

Notes. GAD-7 = Generalized Anxiety Disorder-7; GDS = Geriatric Depression Scale; OR = odds ratio; CI = confidence interval; CBO = community-based organization; SPPB = short physical performance battery.

^a 95% confidence intervals were constructed using cluster-robust standard errors to account for the potential nonindependence of observations within our 11 CBOs and clinics.

of any mental health condition between non-Latinx Whites and non-Latinx Blacks or American Indians, although this could be due to limited power to detect these differences in small samples. In all cases, better physical health (higher SPPB scores) was associated with decreased odds of any mental health condition. Compared to the 60–64 age group, participants ages 75+ had decreased odds of mental health symptoms and suicidal behavior (omnibus tests with $p < .05$). Participants recruited at CBOs had lower odds of *mild-to-severe anxiety* and *depression symptoms* than participants recruited at clinics.

Results from Table 4 indicated significant differences across language groups for *mild-to-severe anxiety* and *depression symptoms* (omnibus test with $p < .05$), but not for *any suicidal behavior*. Individual language coefficients followed similar patterns to race/ethnicity. Compared to those who indicated English as their preferred language, the Spanish language group had higher odds of *mild-to-severe anxiety symptoms* ($OR [95\% CI] = 1.54 [1.02, 2.34]$), while Spanish, Mandarin, and Cantonese language groups all had higher odds of *mild-to-severe depression symptoms* ($OR [95\% CI] = 1.86 [1.20, 2.88]$; $OR [95\% CI] = 2.66 [1.79, 3.96]$; and $OR [95\% CI] = 2.46 [1.71, 3.54]$, respectively). We also observed lower odds of *mild-to-severe anxiety symptoms* for the Mandarin language group ($OR [95\% CI] = .58 [.42, .80]$).

Discussion

Consistent with previous studies reporting higher rates of depressive symptoms among Latinx and Asian older adults compared to the U.S. general population (Glymour et al., 2012; González et al., 2010; Kim et al., 2009; Rote et al., 2015), we found evidence that Latinx and Asian older adults had increased odds of mild-to-severe depression symptoms than non-Latinx Whites. Also consistent with previous studies (Falcón & Tucker, 2000; Shetterly et al., 1998), we found that older Latinx had higher odds of any suicidal behavior than non-Latinx Whites. Higher rates of depressive symptoms and increased risk of suicide in older Latinx have been associated with more medical illness (Briones et al., 2011; Juurlink et al., 2004; Kemp et al., 1987). Consistent with this finding, we found that worse physical health, as measured by lower SPPB scores, was associated with increased odds of mild-to-severe anxiety and depressive symptoms and any suicidal behavior (Patel et al., 2019; Song et al., 2014). The association between physical illnesses and suicide in late life has been partially explained by the mediating effect of depression (Conwell & Thompson, 2008). Older adults who report suicidal behaviors have more depressive illness, physical illness burden, and functional limitations than adults who did not complete suicide (Turvey et al., 2002).

We found that Asians had higher odds of mild-to-severe depressive symptoms than non-Latinx Whites but not of mild-to-severe anxiety symptoms or any suicidal behavior. Higher prevalence of depressive symptoms among Asians has been associated with acculturation stress caused by perceptions of a cultural gap between themselves and their adult children. This cultural gap has been associated with increased concerns about caretaking responsibilities, social isolation, and depression (Mui & Kang, 2006), and worries about access to services (Koehn & Kobayashi, 2011; Nguyen, 2012). This intergenerational tension might explain the high prevalence of depressive symptoms in our sample of Asian older adults.

We hypothesized that adults ages 75+ in all racial/ethnic groups would report less anxiety and depressive symptoms and suicidal behaviors than adults ages 60–74. We found less anxiety and depressive symptoms with increasing age, consistent with U.S. epidemiological evidence (Byers et al., 2010; Hybels & Blazer, 2003). A possible explanation is that adults ages 75+ attending CBOs are likely to be healthier. Our results showed that depression and anxiety symptoms were more prevalent in participants recruited in clinics compared to CBOs. Some studies have also found that older adults are less likely than younger adults to acknowledge symptoms of anxiety and depression (Fisher & Goldney, 2003; Wetherell et al., 2003).

We posited that non-English speakers would report higher levels of mental health symptoms than English-speakers. Indeed, we found that non-English language preference was related to increased anxiety and depression symptoms. Prior research indicates that non-English speaking minority older adults are at a higher risk of social isolation and family disruption (Casado & Leung, 2002; Mui & Kang, 2006). Language preference, a proxy for acculturation, has been linked to symptoms of depression and anxiety (Casado & Leung, 2002; Chiriboga et al., 2002). Latinx who prefer Spanish may have more traditional value structures than bilingual Latinx or who prefer English and may experience social isolation because of language barriers (Santiago-Rivera et al., 2002). Less acculturation and limited English proficiency might limit older Latinx to seek mental health care (Cabassa et al., 2006; González et al., 2001).

Mandarin and Cantonese speaking older adults, all but one who self-identified as Asian, reported significantly more depressive symptoms than English-speakers. However, Mandarin speakers reported less anxiety than English-speaking older adults. Asians are not a homogenous group (Gordon et al., 2019; Mui & Kang, 2006); thus, it is possible that in our sample, Mandarin language preference served as a proxy for less acculturation and was associated with protective family factors which may differ from other older adults with different Asian languages. Prior studies have demonstrated lower rates of anxiety among China-born immigrants who arrived in the U.S. at ages 18 or older compared to those who immigrated prior to age 18 (Zhang et al., 2013). Studies have shown lower rates of anxiety and depression in U.S. Asian older adults than Latinx (Park et al., 2014) and non-Latinx Whites (Kim & López, 2014). This might be related to opportunities to socialize and maintain friendships in their neighborhoods. Family caretaking has a protective effect on depression in foreign-born Asian immigrants, as their children are more likely to assume a caretaker role than children of their American-born counterparts.

As hypothesized, we found that participants from clinics had higher odds of anxiety and depression symptoms than participants from CBOs. This may be partially explained by healthier populations attending CBOs, or perhaps CBOs themselves offering a protective advantage through socialization, routine, and basic needs assistance (Alegría, Frontera, et al., 2019; Siegler et al., 2015).

Our study had several potential limitations. Because we used a convenience sample, the results may not be representative of the general population or generalizable outside the type of organizations examined. The uneven distribution of participants across study locations (most in Massachusetts) also impacts the generalizability of our results. Second, the prevalence obtained in this study reflects minority older adults from CBOs, which may be a healthier population than the average U.S. minority older adult, particularly

those who are homebound. Third, although gender was not significantly associated with the outcomes, we did not have a large enough sample to test differences by race/ethnicity and gender. We also lacked other indicators of socioeconomic status, since the only indicator available was educational level. Given the short screener, we did not collect information about country of birth, years in the U.S., or variables measuring degree of acculturation, family contact frequency, or family arrangement. Language preference was used as a proxy for acculturation, but further research is needed to examine how acculturation may be related to anxiety, depression, and suicide in racial/ethnic older adults in CBOs and clinics. Language preference was also highly correlated with race/ethnicity, which made it impossible to test the association between race/ethnicity and our outcomes controlling for language. For example, 97% of Latinx participants responded to the interview in Spanish, although not all Spanish respondents were Latinx (77% of American Indian and 21% of non-Latinx Black also responded in Spanish). While we found that Latinx older adults had increased odds of any suicidal behavior, this same result did not hold in the Spanish language group. We believe that this was mostly attributable to a very small number of non-Latinx Black and American Indian participants indicating any suicidal behavior. However, we could not disentangle how much of the increased odds in any suicidal behavior among Latinx was explained by ethnicity and how much by language. It is essential for future work to contain large diverse samples to fully evaluate the cultural aspects that contribute to distinct patterns of suicidal behaviors. Data limitations also did not allow us to examine if some study participants used two or more languages or shifted to a current dominant language. Caution should be taken not to generalize by ethnic group, since we were not able to analyze subethnicities. For example, Latinx and Asians are heterogeneous groups with many subethnicities with distinct cultural values and belief systems (Gordon et al., 2019; Jimenez et al., 2019). Finally, no studies assessing measurement invariance by language and race/ethnicity were identified for some of our outcomes; thus, it is possible that the observed differences reflect differences in interpretation rather than differences in mental health outcomes.

Our main finding was the high prevalence of anxiety and depression symptoms and suicidal behaviors in U.S. racial/ethnic minority older adults. The lack of outreach for screening is a barrier to mental healthcare, which might exacerbate health outcome disparity. Undetected mental health symptoms in older minority groups (Aranda et al., 2001) could result in increased morbidity and mortality and negative public health consequences (Unützer et al., 1997), even in the presence of effective treatments to reduce symptoms of anxiety and depression in ethnic minority older adult populations (Alegría, Frontera, et al., 2019).

Conclusion

Our results underscore the importance of outreach and screening for mental health problems in racial/ethnic minority older adults in CBOs and clinics to provide early mental health care, a key measure for preventing the rising burden of mental disorders in older adults. CBOs offer an opportunity to engage racial/ethnic minorities in preventive care to increase awareness of symptoms of mental illness and referral systems for early treatment. CBOs and clinics may be crucial in promoting mental health literacy, engaging patients in care, and preventing mental disorders and greater disability (Jorm,

2012). Clinics can partner with CBOs to serve their patients to promote prevention and wellness. Similarly, CBOs can bridge to clinics, by activating participants to take charge of their health and facilitating the referral process and access to clinics.

References

- Administration for Community Living. (2018). *Profile of older Americans | ACL administration for community living*. <https://acl.gov/aging-and-disability-in-america/data-and-research/profile-older-americans>
- Alegría, M., Frontera, W., Cruz-Gonzalez, M., Markle, S. L., Trinh-Shevrin, C., Wang, Y., Herrera, L., Ishikawa, R. Z., Velazquez, E., Fuentes, L., Guo, Y., Pan, J., Cheung, M., Wong, J., Genatios, U., Jimenez, A., Ramos, Z. I., Perez, G., Wong, J. Y., . . . Shrout, P. E. (2019). Effectiveness of a disability preventive intervention for minority and immigrant elders: The positive minds-strong bodies randomized clinical trial. *The American Journal of Geriatric Psychiatry*, 27(12), 1299–1313. <https://doi.org/10.1016/j.jagp.2019.08.008>
- Almeida, O. P., Draper, B., Snowden, J., Lautenschlager, N. T., Pirkis, J., Byrne, G., Sim, M., Stocks, N., Flicker, L., & Pfaff, J. J. (2012). Factors associated with suicidal thoughts in a large community study of older adults. *The British Journal of Psychiatry*, 201(6), 466–472. <https://doi.org/10.1192/bjp.bp.112.110130>
- Angel, R. J., Frisco, M., Angel, J. L., & Chiriboga, D. A. (2003). Financial strain and health among elderly Mexican-origin individuals. *Journal of Health and Social Behavior*, 44(4), 536–551. <https://doi.org/10.2307/1519798>
- Aranda, M. P., Lee, P. J., & Wilson, S. (2001). Correlates of depression in older Latinos. *Home Health Care Services Quarterly*, 20(1), 1–20. https://doi.org/10.1300/J027v20n01_01
- Bailey, R. K., Mokongho, J., & Kumar, A. (2019). Racial and ethnic differences in depression: Current perspectives. In *Neuropsychiatric disease and treatment*, (Vol. 15, pp. 603–609). Dove Medical Press. <https://doi.org/10.2147/NDT.S128584>
- Balakrishnan, A., & Jordan, N. (2019). The undocumented elderly: Coverage gaps and low health care use. *Journal of Health Care for the Poor and Underserved*, 30(3), 891–898. <https://doi.org/10.1353/hpu.2019.0062>
- Balsamo, M., Cataldi, F., Carlucci, L., Padulo, C., & Fairfield, B. (2018). Assessment of late-life depression via self-report measures: A review. *Clinical Interventions in Aging*, 13, 2021–2044. <https://doi.org/10.2147/CIA.S114100>
- Bennardi, M., Caballero, F. F., Miret, M., Ayuso-Mateos, J. L., Haro, J. M., Lara, E., Arensman, E., & Cabello, M. (2019). Longitudinal relationships between positive affect, loneliness, and suicide ideation: Age-specific factors in a general population. *Suicide & Life-Threatening Behavior*, 49(1), 90–103. <https://doi.org/10.1111/sltb.12424>
- Blas, E., Gilson, L., Kelly, M. P., Labonté, R., Lapitan, J., Muntaner, C., Östlin, P., Popay, J., Sadana, R., Sen, G., Schrecker, T., & Vaghri, Z. (2008). Addressing social determinants of health inequities: What can the state and civil society do? *Lancet*, 372(9650), 1684–1689. [https://doi.org/10.1016/S0140-6736\(08\)61693-1](https://doi.org/10.1016/S0140-6736(08)61693-1)
- Briones, D. F., Heller, P. L., Carcoba, L. M., Weisman, H. W., Ledger, E. M., & Escamilla, M. A. (2011). Health-related conditions and depression in elderly Mexican American and non-Hispanic white residents of a United States-Mexico border county: Moderating effects of educational attainment. *Depression Research and Treatment*, 2011, Article 908536. <https://doi.org/10.1155/2011/908536>
- Byers, A. L., Yaffe, K., Covinsky, K. E., Friedman, M. B., & Bruce, M. L. (2010). High occurrence of mood and anxiety disorders among older adults: The National Comorbidity Survey replication. *Archives of General Psychiatry*, 67(5), 489–496. <https://doi.org/10.1001/archgenpsychiatry.2010.35>
- Cabassa, L. J., Zayas, L. H., & Hansen, M. C. (2006). Latino adults' access to mental health care: A review of epidemiological studies. *Administration*

- and Policy in Mental Health, 33(3), 316–330. <https://doi.org/10.1007/s10488-006-0040-8>
- Caplan, S. (2007). Latinos, acculturation, and acculturative stress: A dimensional concept analysis. *Policy, Politics & Nursing Practice*, 8(2), 93–106. <https://doi.org/10.1177/1527154407301751>
- Casado, B. L., & Leung, P. (2002). Migratory grief and depression among elderly Chinese American immigrants. *Journal of Gerontological Social Work*, 36(1–2), 5–26. https://doi.org/10.1300/J083v36n01_02
- Cassidy, S. A., Bradley, L., Bowen, E., Wigham, S., & Rodgers, J. (2018). Measurement properties of tools used to assess suicidality in autistic and general population adults: A systematic review. *Clinical Psychology Review*, 62(3), 56–70. <https://doi.org/10.1016/j.cpr.2018.05.002>
- Chang, Q., Chan, C. H., & Yip, P. S. F. (2017). A meta-analytic review on social relationships and suicidal ideation among older adults. *Social Science & Medicine*, 191, 65–76. <https://doi.org/10.1016/j.socscimed.2017.09.003>
- Chen, W. Q., Siu, O. L., Lu, J. F., Cooper, C. L., & Phillips, D. R. (2009). Work stress and depression: The direct and moderating effects of informal social support and coping. *Stress and Health*, 25(5), 431–443. <https://doi.org/10.1002/smi.1263>
- Chillag, K., Bartholow, K., Cordeiro, J., Swanson, S., Patterson, J., Stebbins, S., Woodside, C., & Sy, F. (2002). Factors affecting the delivery of HIV/AIDS prevention programs by community-based organizations. *AIDS Education and Prevention*, 14(3, Suppl A), 27–37. <https://doi.org/10.1521/aeap.14.4.27.23886>
- Chiriboga, D. A., Black, S. A., Aranda, M., & Markides, K. (2002). Stress and depressive symptoms among Mexican American elders. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 57(6), 559–568. <https://doi.org/10.1093/geronb/57.6.P559>
- Cokley, K., & Moore, P. (2007). Moderating and mediating effects of gender and psychological disengagement on the academic achievement of African American college students. *The Journal of Black Psychology*, 33(2), 169–187. <https://doi.org/10.1177/0095798407299512>
- Conwell, Y., & Thompson, C. (2008). Suicidal behavior in elders. *The Psychiatric Clinics of North America*, 31(2), 333–356. <https://doi.org/10.1016/j.psc.2008.01.004>
- Cullen, P. (2019). Mental illness in the oldest-old. In *Advanced age geriatric care* (pp. 145–157). Springer International Publishing. https://doi.org/10.1007/978-3-319-96998-5_17
- Davison, K. M., Lung, Y., Lin, S. L., Tong, H., Kobayashi, K. M., & Fuller-Thomson, E. (2019). Depression in middle and older adulthood: The role of immigration, nutrition, and other determinants of health in the Canadian longitudinal study on aging. *BMC Psychiatry*, 19(1), Article 329. <https://doi.org/10.1186/s12888-019-2309-y>
- Derr, A. S. (2016). Mental health service use among immigrants in the United States: A systematic review. *Psychiatric Services*, 67(3), 265–274. <https://doi.org/10.1176/appi.ps.201500004>
- DuBard, C. A., & Gizlice, Z. (2008). Language spoken and differences in health status, access to care, and receipt of preventive services among US Hispanics. *American Journal of Public Health*, 98(11), 2021–2028. <https://doi.org/10.2105/AJPH.2007.119008>
- Ell, K. (2006). Depression care for the elderly: Reducing barriers to evidence-based practice. *Home Health Care Services Quarterly*, 25(1–2), 115–148. https://doi.org/10.1300/J027v25n01_07
- Evans-Lacko, S., Corker, E., Williams, P., Henderson, C., & Thornicroft, G. (2014). Effect of the Time to Change anti-stigma campaign on trends in mental-illness-related public stigma among the English population in 2003–13: An analysis of survey data. *The Lancet. Psychiatry*, 1(2), 121–128. [https://doi.org/10.1016/S2215-0366\(14\)70243-3](https://doi.org/10.1016/S2215-0366(14)70243-3)
- Falcón, L. M., & Tucker, K. L. (2000). Prevalence and correlates of depressive symptoms among Hispanic elders in Massachusetts. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 55(2), S108–S116. <https://doi.org/10.1093/geronb/55.2.S108>
- Fässberg, M. M., Cheung, G., Canetto, S. S., Erlangsen, A., Lapierre, S., Lindner, R., Draper, B., Gallo, J. J., Wong, C., Wu, J., Duberstein, P., & Wærn, M. (2016). A systematic review of physical illness, functional disability, and suicidal behaviour among older adults. *Aging & Mental Health*, 20(2), 166–194. <https://doi.org/10.1080/13607863.2015.1083945>
- Feng, L., Yap, K. B., & Ng, T. P. (2013). Depressive symptoms in older adults with chronic kidney disease: Mortality, quality of life outcomes, and correlates. *The American Journal of Geriatric Psychiatry*, 21(6), 570–579. <https://doi.org/10.1016/j.jagp.2012.12.020>
- Fisher, L. J., & Goldney, R. D. (2003). Differences in community mental health literacy in older and younger Australians. *International Journal of Geriatric Psychiatry*, 18(1), 33–40. <https://doi.org/10.1002/gps.769>
- García, C., García, M. A., Chiu, C. T., Rivera, F. I., & Raji, M. (2019). Life expectancies with depression by age of migration and gender among older mexican americans. *The Gerontologist*, 59(5), 877–885. <https://doi.org/10.1093/geront/gny107>
- Gerst, K., Al-Ghatrif, M., Beard, H. A., Samper-Ternent, R., & Markides, K. S. (2010). High depressive symptomatology among older community-dwelling Mexican Americans: The impact of immigration. *Aging & Mental Health*, 14(3), 347–354. <https://doi.org/10.1080/13607860903292578>
- Glover, L. M., Sims, M., & Winters, K. (2017). Perceived discrimination and reported trust and satisfaction with providers in African Americans: The Jackson heart study. *Ethnicity & Disease*, 27(3), 209–216. <https://doi.org/10.18865/ed.27.3.209>
- Glymour, M. M., Yen, J. J., Kosheleva, A., Moon, J. R., Capistrant, B. D., & Patton, K. K. (2012). Elevated depressive symptoms and incident stroke in Hispanic, African-American, and White older Americans. *Journal of Behavioral Medicine*, 35(2), 211–220. <https://doi.org/10.1007/s10865-011-9356-2>
- González, H. M., Haan, M. N., & Hinton, L. (2001). Acculturation and the prevalence of depression in older Mexican Americans: Baseline results of the Sacramento Area Latino study on aging. *Journal of the American Geriatrics Society*, 49(7), 948–953. <https://doi.org/10.1046/j.1532-5415.2001.49186.x>
- González, H. M., Tarraf, W., Whitfield, K. E., & Vega, W. A. (2010). The epidemiology of major depression and ethnicity in the United States. *Journal of Psychiatric Research*, 44(15), 1043–1051. <https://doi.org/10.1016/j.jpsychires.2010.03.017>
- Gordon, N. P., Lin, T. Y., Rau, J., & Lo, J. C. (2019). Aggregation of Asian-American subgroups masks meaningful differences in health and health risks among Asian ethnicities: An electronic health record based cohort study. *BMC Public Health*, 19(1), Article 1551. <https://doi.org/10.1186/s12889-019-7683-3>
- Grenier, S., Payette, M. C., Gunther, B., Askari, S., Desjardins, F. F., Raymond, B., & Berbiche, D. (2019). Association of age and gender with anxiety disorders in older adults: A systematic review and meta-analysis. *International Journal of Geriatric Psychiatry*, 34(3), 397–407. <https://doi.org/10.1002/gps.5035>
- Guralnik, J. M., Simonsick, E. M., Ferrucci, L., Glynn, R. J., Berkman, L. F., Blazer, D. G., Scherr, P. A., & Wallace, R. B. (1994). A short physical performance battery assessing lower extremity function: Association with self-reported disability and prediction of mortality and nursing home admission. *Journal of Gerontology*, 49(2), M85–M94. <https://doi.org/10.1093/geronj/49.2.M85>
- Hansen, B. R., Hodgson, N. A., & Gitlin, L. N. (2016). It's a matter of trust: Older African Americans speak about their health care encounters. *Journal of Applied Gerontology*, 35(10), 1058–1076. <https://doi.org/10.1177/0733464815570662>
- Hegeman, J. M., Kok, R. M., van der Mast, R. C., & Giltay, E. J. (2012). Phenomenology of depression in older compared with younger adults: Meta-analysis. *The British Journal of Psychiatry*, 200(4), 275–281. <https://doi.org/10.1192/bjp.bp.111.095950>
- Ho, C. Sh., Feng, L., Fam, J., Mahendran, R., Kua, E. H., & Ng, T. P. (2014). Coexisting medical comorbidity and depression: Multiplicative effects on

- health outcomes in older adults. *International Psychogeriatrics*, 26(7), 1221–1229. <https://doi.org/10.1017/S1041610214000611>
- Holden, K. B., Bradford, L. D., Hall, S. P., & Belton, A. S. (2013). Prevalence and correlates of depressive symptoms and resiliency among African American women in a community-based primary health care center. *Journal of Health Care for the Poor and Underserved*, 24(4, Suppl), 79–93. <https://doi.org/10.1353/hpu.2014.0012>
- Hong, S., Walton, E., Tamaki, E., & Sabin, J. A. (2014). Lifetime prevalence of mental disorders among Asian Americans: Nativity, gender, and socio-demographic correlates. *Asian American Journal of Psychology*, 5(4), 353–363. <https://doi.org/10.1037/a0035680>
- Hybels, C. F., & Blazer, D. G. (2003). Epidemiology of late-life mental disorders. *Clinics in Geriatric Medicine*, 19(4), 663–696. [https://doi.org/10.1016/S0749-0690\(03\)00042-9](https://doi.org/10.1016/S0749-0690(03)00042-9)
- Jeste, D. V., Palmer, B. W., Appelbaum, P. S., Golshan, S., Glorioso, D., Dunn, L. B., Kim, K., Meeks, T., & Kraemer, H. C. (2007). A new brief instrument for assessing decisional capacity for clinical research. *Archives of General Psychiatry*, 64(8), 966–974. <https://doi.org/10.1001/archpsyc.64.8.966>
- Jimenez, A., Alegria, M., Camino, R., Lazaro, J., & Lisotto, M. J. (2019). Cultural sensitivity: what should we understand about Latinos? In R. Parekh (Ed.), *The Massachusetts general hospital textbook on diversity and cultural sensitivity in mental health, current clinical psychiatry* (2nd ed., pp. 201–228). Springer. https://doi.org/10.1007/978-3-030-20174-6_11
- Jimenez, D. E., Bartels, S. J., Cardenas, V., & Alegria, M. (2013). Stigmatizing attitudes toward mental illness among racial/ethnic older adults in primary care. *International Journal of Geriatric Psychiatry*, 28(10), 1061–1068. <https://doi.org/10.1002/gps.3928>
- Jimenez, D. E., Bartels, S. J., Cardenas, V., Dhaliwal, S. S., & Alegria, M. (2012). Cultural beliefs and mental health treatment preferences of ethnically diverse older adult consumers in primary care. *The American Journal of Geriatric Psychiatry*, 20(6), 533–542. <https://doi.org/10.1097/JGP.0b013e318227f876>
- Jones, A. (2012). Disability, health and generation status: How Hispanics in the US fare in late life. *Journal of Immigrant and Minority Health*, 14(3), 467–474. <https://doi.org/10.1007/s10903-011-9500-7>
- Jones, A. L., Cochran, S. D., Leibowitz, A., Wells, K. B., Kominski, G., & Mays, V. M. (2018). Racial, ethnic, and nativity differences in mental health visits to primary care and specialty mental health providers: Analysis of the medical expenditures panel survey, 2010–2015. *Health-care*, 6(2), Article 29. <https://doi.org/10.3390/healthcare6020029>
- Jorm, A. F. (2012). Mental health literacy: Empowering the community to take action for better mental health. *American Psychologist*, 67(3), 231–243. <https://doi.org/10.1037/a0025957>
- Juurlink, D. N., Herrmann, N., Szalai, J. P., Kopp, A., & Redelmeier, D. A. (2004). Medical illness and the risk of suicide in the elderly. *Archives of Internal Medicine*, 164(11), 1179–1184. <https://doi.org/10.1001/archinte.164.11.1179>
- Kahn, J. R., & Fazio, E. M. (2005). Economic status over the life course and racial disparities in health. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 60, 76–84. https://doi.org/10.1093/geronb/60.special_issue_2.s76
- Kang, S. Y., Basham, R., & Kim, Y. J. (2013). Contributing factors of depressive symptoms among elderly Korean immigrants in Texas. *Journal of Gerontological Social Work*, 56(1), 67–82. <https://doi.org/10.1080/01634372.2012.734369>
- Kemp, B. J., Staples, F., & Lopez-Aqueres, W. (1987). Epidemiology of depression and dysphoria in an elderly Hispanic population. Prevalence and correlates. *Journal of the American Geriatrics Society*, 35(10), 920–926. <https://doi.org/10.1111/j.1532-5415.1987.tb02293.x>
- Kim, G., Chiriboga, D. A., & Jang, Y. (2009). Cultural equivalence in depressive symptoms in older white, black, and Mexican-American adults. *Journal of the American Geriatrics Society*, 57(5), 790–796. <https://doi.org/10.1111/j.1532-5415.2009.02188.x>
- Kim, G., Kim, M., Park, S., Jimenez, D. E., & Chiriboga, D. A. (2019). Limited English proficiency and trajectories of depressive symptoms among Mexican American older adults. *The Gerontologist*, 59(5), 856–864. <https://doi.org/10.1093/geront/gny032>
- Kim, G., Worley, C. B., Allen, R. S., Vinson, L., Crowther, M. R., Parmelee, P., & Chiriboga, D. A. (2011). Vulnerability of older Latino and Asian immigrants with limited English proficiency. *Journal of the American Geriatrics Society*, 59(7), 1246–1252. <https://doi.org/10.1111/j.1532-5415.2011.03483.x>
- Kim, H. J., Park, E., Storr, C. L., Tran, K., & Juon, H. S. (2015). Depression among Asian-American adults in the community: Systematic review and meta-analysis. *PLOS ONE*, 10(6), Article e0127760. <https://doi.org/10.1371/journal.pone.0127760>
- Kim, J. M., & López, S. R. (2014). The expression of depression in Asian Americans and European Americans. *Journal of Abnormal Psychology*, 123(4), 754–763. <https://doi.org/10.1037/a0038114>
- Kiosses, D. N., & Marino, P. (2019). Assessment of depression and anxiety in older adults. In L. Ravdin & H. Katzen (Eds.), *Handbook on the neuropsychology of aging and dementia* (pp. 49–59). Springer. https://doi.org/10.1007/978-3-319-93497-6_4
- Knight, J. (2019). Understanding and managing depression in older people. *Nursing Older People*, 31(6), 41–48. <https://doi.org/10.7748/nop.2019.e1138>
- Koehn, S., & Kobayashi, K. (2011). Age and ethnicity. In *Age discrimination and diversity: Multiple discrimination from an age perspective* (pp. 132–159). Cambridge University Press. <https://doi.org/10.1017/CBO9780511777196.007>
- Krause, N. (1987). Chronic financial strain, social support, and depressive symptoms among older adults. *Psychology and Aging*, 2(2), 185–192. <https://doi.org/10.1037/0882-7974.2.2.185>
- Krause, N., Jay, G., & Liang, J. (1991). Financial strain and psychological well-being among the American and Japanese elderly. *Psychology and Aging*, 6(2), 170–181. <https://doi.org/10.1037/0882-7974.6.2.170>
- Kroenke, K., Spitzer, R. L., Williams, J. B. W., Monahan, P. O., & Löwe, B. (2007). Anxiety disorders in primary care: Prevalence, impairment, comorbidity, and detection. *Annals of Internal Medicine*, 146(5), 317–325. <https://doi.org/10.7326/0003-4819-146-5-200703060-00004>
- Laborde-Lahoz, P., El-Gabalawy, R., Kinley, J., Kirwin, P. D., Sareen, J., & Pietrzak, R. H. (2015). Subsyndromal depression among older adults in the USA: Prevalence, comorbidity, and risk for new-onset psychiatric disorders in late life. *International Journal of Geriatric Psychiatry*, 30(7), 677–685. <https://doi.org/10.1002/gps.4204>
- Lau, A. S., McCabe, K. M., Yeh, M., Garland, A. F., Wood, P. A., & Hough, R. L. (2005). The acculturation gap-distress hypothesis among high-risk Mexican American families. *Journal of Family Psychology*, 19(3), 367–375. <https://doi.org/10.1037/0893-3200.19.3.367>
- Lum, T. Y., & Vanderaa, J. P. (2010). Health disparities among immigrant and non-immigrant elders: The association of acculturation and education. *Journal of Immigrant and Minority Health*, 12(5), 743–753. <https://doi.org/10.1007/s10903-008-9225-4>
- Lund, C., Brooke-Sumner, C., Baingana, F., Baron, E. C., Breuer, E., Chandra, P., Haushofer, J., Herrman, H., Jordans, M., Kieling, C., Medina-Mora, M. E., Morgan, E., Omigbodun, O., Tol, W., Patel, V., & Saxena, S. (2018). Social determinants of mental disorders and the sustainable development goals: A systematic review of reviews. *The Lancet. Psychiatry*, 5(4), 357–369. [https://doi.org/10.1016/S2215-0366\(18\)30060-9](https://doi.org/10.1016/S2215-0366(18)30060-9)
- Luoma, J. B., Martin, C. E., & Pearson, J. L. (2002). Contact with mental health and primary care providers before suicide: A review of the evidence. *The American Journal of Psychiatry*, 159(6), 909–916. <https://doi.org/10.1176/appi.ajp.159.6.909>
- Lutz, J., & Fiske, A. (2018). Functional disability and suicidal behavior in middle-aged and older adults: A systematic critical review. *Journal of Affective Disorders*, 227(6), 260–271. <https://doi.org/10.1016/j.jad.2017.10.043>

- Medina, L. D., Sabo, S., & Vespa, J. (2020). *Living longer: Historical and projected life expectancy in the United States, 1960 to 2060 Current Population Reports, Issue*. <https://www.census.gov/library/publications/2020/demo/p25-1145.html>
- Mills, S. D., Fox, R. S., Malcarne, V. L., Roesch, S. C., Champagne, B. R., & Sadler, G. R. (2014). The psychometric properties of the generalized anxiety disorder-7 scale in Hispanic Americans with English or Spanish language preference. *Cultural Diversity & Ethnic Minority Psychology*, 20(3), 463–468. <https://doi.org/10.1037/a0036523>
- Morin, R. T., Nelson, C., Bickford, D., Insel, P. S., & Mackin, R. S. (2020). Somatic and anxiety symptoms of depression are associated with disability in late life depression. *Aging & Mental Health*, 24(8), 1225–1228. <https://doi.org/10.1080/13607863.2019.1597013>
- Mui, A. C., & Kang, S. Y. (2006). Acculturation stress and depression among Asian immigrant elders. *Social Work*, 51(3), 243–255. <https://doi.org/10.1093/sw/51.3.243>
- Musa, D., Schulz, R., Harris, R., Silverman, M., & Thomas, S. B. (2009). Trust in the health care system and the use of preventive health services by older black and white adults. *American Journal of Public Health*, 99(7), 1293–1299. <https://doi.org/10.2105/AJPH.2007.123927>
- Nguyen, D. (2012). The effects of sociocultural factors on older Asian Americans' access to care. *Journal of Gerontological Social Work*, 55(1), 55–71. <https://doi.org/10.1080/01634372.2011.618525>
- Park, M., & Unützer, J. (2011). Geriatric depression in primary care. *The Psychiatric Clinics of North America*, 34(2), 469–487, ix–x. <https://doi.org/10.1016/j.psc.2011.02.009>
- Park, M., Unützer, J., & Grembowski, D. (2014). Ethnic and gender variations in the associations between family cohesion, family conflict, and depression in older Asian and Latino adults. *Journal of Immigrant and Minority Health*, 16(6), 1103–1110. <https://doi.org/10.1007/s10903-013-9926-1>
- Patel, K. V., Guralnik, J. M., Phelan, E. A., Gell, N. M., Wallace, R. B., Sullivan, M. D., & Turk, D. C. (2019). Symptom burden among community-dwelling older adults in the United States. *Journal of the American Geriatrics Society*, 67(2), 223–231. <https://doi.org/10.1111/jgs.15673>
- Paykel, E. S., Myers, J. K., Lindenthal, J. J., & Tanner, J. (1974). Suicidal feelings in the general population: A prevalence study. *The British Journal of Psychiatry*, 124(582), 460–469. <https://doi.org/10.1192/bjp.124.5.460>
- Perreira, K. M., Gotman, N., Isasi, C. R., Arguelles, W., Castañeda, S. F., Daviglus, M. L., Giachello, A. L., Gonzalez, P., Penedo, F. J., Salgado, H., & Wassertheil-Smoller, S. (2015). Mental health and exposure to the United States: Key correlates from the hispanic community health study of Latinos. *Journal of Nervous and Mental Disease*, 203(9), 670–678. <https://doi.org/10.1097/NMD.0000000000000350>
- Ramanadhan, S., Crisostomo, J., Alexander-Molloy, J., Gandelman, E., Grullon, M., Lora, V., Reeves, C., Savage, C., Viswanath, K., & the PLANET MassCONNECT C-PAC. (2012). Perceptions of evidence-based programs among community-based organizations tackling health disparities: A qualitative study. *Health Education Research*, 27(4), 717–728. <https://doi.org/10.1093/her/cyr088>
- Richardson, T. M., Simming, A., He, H., & Conwell, Y. (2011). Anxiety and its correlates among older adults accessing aging services. *International Journal of Geriatric Psychiatry*, 26(1), 31–38. <https://doi.org/10.1002/gps.2474>
- Rote, S., Chen, N. W., & Markides, K. (2015). Trajectories of depressive symptoms in elderly Mexican Americans. *Journal of the American Geriatrics Society*, 63(7), 1324–1330. <https://doi.org/10.1111/jgs.13480>
- Sadavoy, J., Meier, R., & Ong, A. Y. M. (2004). Barriers to access to mental health services for ethnic seniors: The Toronto study. *Canadian Journal of Psychiatry*, 49(3), 192–199. <https://doi.org/10.1177/070674370404900307>
- Santiago-Rivera, A. L., Arredondo, P., & Gallardo-Cooper, M. (2002). *Counseling Latinos and la Familia: A Practical Guide*. Sage Publications. https://books.google.com.pr/books?hl=es&lr=&id=7Uc5DQAAQBAJ&oi=fnd&pg=PP1&dq=Santiago-Rivera+et+al.,+2002&ots=wki2leDac&sig=hwgh6zqo_JcYyHUd-6UNr9GUK1s&redir_esc=y#v=onepage&q=Santiago-Rivera%20et%20al.%2C%202002&f=false
- Sentell, T., Shumway, M., & Snowden, L. (2007). Access to mental health treatment by English language proficiency and race/ethnicity. *Journal of General Internal Medicine*, 22(2, Suppl 2), 289–293. <https://doi.org/10.1007/s11606-007-0345-7>
- Sheikh, J. I., & Yesavage, J. A. (1986). 9/geriatric depression scale (Gds) recent evidence and development of a shorter version. *Clinical Gerontologist*, 5(1–2), 165–173. https://doi.org/10.1300/J018v05n01_09
- Shetterly, S. M., Baxter, J., Morgenstern, N. E., Grigsby, J., & Hamman, R. F. (1998). Higher instrumental activities of daily living disability in Hispanics compared with non-Hispanic whites in rural Colorado. The San Luis Valley health and aging study. *American Journal of Epidemiology*, 147(11), 1019–1027. <https://doi.org/10.1093/oxfordjournals.aje.a009395>
- Siegler, E. L., Lama, S. D., Knight, M. G., Laureano, E., & Reid, M. C. (2015). Community-based supports and services for older adults: A primer for clinicians. *Journal of Geriatrics*, 2015, 1–6. <https://doi.org/10.1155/2015/678625>
- Snowden, L. R., & Pingitore, D. (2002). Frequency and scope of mental health service delivery to African Americans in primary care. *Mental Health Services Research*, 4(3), 123–130. <https://doi.org/10.1023/A:1019709728333>
- Solar, O., & Irwin, A. (2010). *A conceptual framework for action on the social determinants of health*. World Health Organization. <https://apps.who.int/iris/handle/10665/44489>
- Song, H. J., Meade, K., Akobundu, U., & Sahyoun, N. R. (2014). Depression as a correlate of functional status of community-dwelling older adults: Utilizing a short-version of 5-item Geriatric Depression Scale as a screening tool. *The Journal of Nutrition, Health & Aging*, 18(8), 765–770. <https://doi.org/10.1007/s12603-014-0542-0>
- Spitzer, R. L., Kroenke, K., Williams, J. B. W., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: The GAD-7. *Archives of Internal Medicine*, 166(10), 1092–1097. <https://doi.org/10.1001/archinte.166.10.1092>
- Stata Corp. (2017). *Stata* | StataCorp LLC. <https://www.stata.com/company/>
- Tai-Seale, M., McGuire, T., Colenda, C., Rosen, D., & Cook, M. A. (2007). Two-minute mental health care for elderly patients: Inside primary care visits. *Journal of the American Geriatrics Society*, 55(12), 1903–1911. <https://doi.org/10.1111/j.1532-5415.2007.01467.x>
- Torres, L., Bird, C., & Mata-Greve, F. (2020). Culture and ethnicity in the mental health of older adults. In N. Hantke, A. Etkin, & R. O'Hara (Eds.), *Handbook of mental health and aging* (pp. 23–36). Elsevier. <https://doi.org/10.1016/B978-0-12-800136-3.00003-X>
- Turner, J. J., Adams-Price, C. E., & Wilmoth, J. D. (2019). Service need and older adults 142. *Journal of Human Sciences and Extension*, 7(3), 142–157. <https://www.jhseonline.com/article/view/908>
- Turvey, C. L., Conwell, Y., Jones, M. P., Phillips, C., Simonsick, E., Pearson, J. L., & Wallace, R. (2002). Risk factors for late-life suicide: A prospective, community-based study. *The American Journal of Geriatric Psychiatry*, 10(4), 398–406. <https://doi.org/10.1097/00019442-200207000-00006>
- Unützer, J., Patrick, D. L., Simon, G., Grembowski, D., Walker, E., Rutter, C., & Katon, W. (1997). Depressive symptoms and the cost of health services in HMO patients aged 65 years and older. A 4-year prospective study. *Journal of the American Medical Association*, 277(20), 1618–1623. <https://doi.org/10.1001/jama.1997.03540440052032>
- Van Orden, K. A., & Conwell, Y. (2016). Issues in research on aging and suicide. *Aging & Mental Health*, 20(2), 240–251. <https://doi.org/10.1080/13607863.2015.1065791>
- Ward, M. C., Miller, B. F., Marconi, V. C., Kaslow, N. J., & Farber, E. W. (2016). The role of behavioral health in optimizing care for complex patients in the primary care setting. *Journal of General Internal Medicine*, 31(3), 265–267. <https://doi.org/10.1007/s11606-015-3499-8>

- Wetherell, J. L., Gatz, M., & Craske, M. G. (2003). Treatment of generalized anxiety disorder in older adults. *Journal of Consulting and Clinical Psychology, 71*(1), 31–40. <https://doi.org/10.1037/0022-006X.71.1.31>
- Wisnivesky, J. P., Krauskopf, K., Wolf, M. S., Wilson, E. A. H., Sofianou, A., Martynenko, M., Halm, E. A., Leventhal, H., & Federman, A. D. (2012). The association between language proficiency and outcomes of elderly patients with asthma. *Annals of Allergy, Asthma & Immunology, 109*(3), 179–184. <https://doi.org/10.1016/j.anai.2012.06.016>
- World Health Organization. (2014). *Social determinants of mental health*. <https://apps.who.int/iris/bitstream/handle/10665/112828/9789?sequence=1>
- World Health Organization. (2021). *Global report on ageism*. <http://www.who.int/publications/i/item/9789240016866>
- Wu, A. W., Weston, C. M., Ibe, C. A., Ruberman, C. F., Bone, L., Boonyasai, R. T., Hwang, S., Gentry, J., Purnell, L., Lu, Y., Liang, S., & Rosenblum, M. (2019). The Baltimore community-based organizations neighborhood network: Enhancing capacity together (CONNECT) cluster RCT. *American Journal of Preventive Medicine, 57*(2), e31–e41. <https://doi.org/10.1016/j.amepre.2019.03.013>
- Yang, F., & Levkoff, S. (2005). Ageism and minority populations: Strengths in the face of challenge. *Generations, 29*(3), 42–48.
- Zhang, J., Fang, L., Wu, Y. W. B., & Wiecezorek, W. F. (2013). Depression, anxiety, and suicidal ideation among Chinese Americans: A study of immigration-related factors. *Journal of Nervous and Mental Disease, 201*(1), 17–22. <https://doi.org/10.1097/NMD.0b013e31827ab2e2>

Received July 2, 2020

Revision received October 14, 2021

Accepted December 19, 2021 ■